(oral drench)

Product Profile

Give every pig a fighting chance.

Nutritional support for newborn, fall back and at-risk piglets.

Key Benefits

- Nutritionally supports piglets with scours in health-challenged situations
- Highly effective for rescuing fall back pigs
- Can be used as a nutritional supplement when quality colostrum is limited.
- Supports health, milk intake and nutrient absorption
- Promotes:
 - Livability
 - Number of full-value pigs weaned
 - · Greater weaning weight
 - Uniformity in weight of weaned pigs
- Supports gut health

Product Description

 An oral drench consisting of a unique and proprietary combination of probiotic bacteria, concentrated yeastbased fermentation extract, micronutrients, pathogenspecific spray-dried polyclonal egg antibodies, organic acids and botanical extracts.

Label Indications & Mixing Directions

- No label restrictions
- 1 dose equals one 2 ml pump
- Standard usage:
 - Supplement all piglets 12 to 36 hours after birth with one dose. Repeat after 24 hours as needed.
- Use with large litters:
 - Give to the smallest 1/3 of piglets as nutritional support. Repeat after 24 hours as needed.
- Use during health challenges:
 - Give one dose to pigs in fall back pens in nurseries and wean-to-finish barns. Repeat after 24 hours as needed.

Packaging

- 6-16 ounce bottles (1,200 doses) with 2-2 ml pump
- 12 8 ounce bottles (1,200 doses) with 12 2 ml pumps

Product Formulation

Thick, brownish liquid with a strong aroma

Regulatory Status

- Natural feed supplement that uses only AAFCO-approved ingredients
- No Veterinary Feed Directive (VFD) required
- Not for human consumption

Storage & Handling

- Store in a cool, dry place
- Shake well before use
- Avoid contact with skin and eyes
- Avoid freezing

Note: Always review Safety Data Sheet (SDS) prior to handling.



Product Codes:

FMC400

6 - 16 oz bottles + 2 drenchers

FMC420

12 - 8 oz bottles + 12 drenchers



Furst Boost™

Standard Operating Procedure



Step 1

Inspect bottles to ensure all are intact.



Step 2

Store in a cool, dry place. Keep out of reach of children.



Step 3

Prior to use, put on appropriate Personal Protective Equipment (PPE), which may include, but not limited to:

- Eye/face protection
 - · Safety glasses or chemical goggles
- Skin protection

Note: Review the Safety Data Sheet (SDS) before product use



Step 4

Determine which piglets will receive Furst Boost™.

- Shake well before use
- Properly secure and support the piglet
- Properly place the dosing tip in the piglet's mouth at the side of the cheek
 - Do not apply directly into the piglet's throat to avoid accidental aspiration into the piglet's respiratory tract
- Proceed to gently depress the plunger and completely dispense the full dose of Furst Boost™
- Dosage:
 - 1 dose = one 2ml pump
 - Standard Usage: Administer 1 dose 12 to 36 hours after birth. Repeat with 2nd dose 24 hours later as needed.
 - Large litters: Administer 1 dose to the smallest 1/3 of the piglets. Repeat with 2nd dose 24 hours later as needed.
 - Health challenges: Administer 1 dose to pigs in fall back pens in nurseries and wean-to-finish barns. Repeat with 2nd dose 24 hours later as needed.

