

FURST FORCE™ CA

SUPPORT YOUR CALVES DURING STRESSFUL SITUATIONS

Weather, socialization, dystocia, colostrum and milk intake, environmental pathogens can all create issues for calves less able to withstand such challenges.

FURST FORCE™ CALF is a unique oral drench for newborn calves utilizing a complex blend of nutritional technologies designed to support the animal, while providing the gastro-intestinal support calves need, especially those at-risk.

FURST FORCE™ CALF delivers:

- 1. Three specific strains of probiotic bacteria (B. pumilus, B. subtilis and B. licheniformis) with a concentrated yeast-based fermentation extract that contains many micro-nutrients to support gut health.
- 2. A specific blend of spray dried egg powder that is a source of polyclonal egg antibodies to **support** health and immunity.
- 3. Specific organic acids to **support digestion** as well as product stability.
- 4. Beta Glucans and MOS for added health benefits.
- 5. Vegetable fat as an added energy source.
- Key botanical extracts, essential oils and flavorings to support health, immunity and stimulate suckling.

Key responses that have been observed



Reduced **Death Loss**



Decreased Scours






Reduced **Digestive Upsets**



Improved Calf Viability



Boosted IgA Titers



FURST FORCE™ CALF

LIFETIME HEALTH AND PRODUCTION STARTS WITH FURST FORCETM CALF



Guaranteed Analysis	
Crude Protein, min.	9.30%
Crude Fat, min.	8.00%
Crude Fiber, max.	3.50%
Calcium (Ca), min.	0.01%
Calcium (Ca), max.	0.09%
Phosphorus (P), min.	0.15%
Potassium (K), min.	0.25%

Ingredients:
Propylene Glycol, Water, Molasses Products, Animal Protein Products, Vegetable Fat, Plant
Protein Products, Saccharomyces Cerevisiae Fermentation Extract, Acetic Acid, Bentonite,
Hydrated Sodium Calcium Aluminosilicate, Lactic Acid, Spices and Other Natural Seasonings and Flavorings.

Directions for Use:BEEF: Give a 15 mL dose orally from birth to 72 hours to supplement calves, especially smaller, at-risk calves.

DAIRY: Give a 15 mL dose orally as soon after colostrum supplementation as practicable. Can be followed by a second dose 12 hours later if necessary.

OTHER: Give a 10 mL dose orally at the onset of scours to support the nutritional needs of the calf. Continue at the same rate for 2-3 consecutive days if necessary.

Individual Bottle Code: FMC415

16 oz. (473 mL) bottle

Case Code: FMC404

6 x 16 oz. (473 mL) bottles (2,838 mL/case)

ACTUAL SIZE!